

AMENDMENTS TO THE CLAIMS


Please amend claims 1, 26, 56-57, 60, 62, 63, 69, 81, and 86 and please cancel claims 2-25, 27-55, 58, 59, 61, 64, 67, 68, 71, 74, 80, and 82-85 without prejudice or disclaimer. The following listing of the claims will replace all prior versions, and listings, of claims in the application.

1. (Currently amended) A method of inhibiting the GTPase activity of dynamin in a cell or synaptosome, comprising contacting a cell or synaptosome with an effective amount of a compound of formula I, or a physiologically acceptable salt thereof, to inhibit said GTPase activity in said cell or synaptosome, wherein



Formula I

M and M' are each independently a moiety of formula II and are the same or different, and Sp is a spacer comprising a 1 to 7 atom chain;



Formula II

V is C or CH;

W is CH or a linker group of up to 3 atoms in length; and

Y is cyano, nitro, NH, amino, oxo, halo, hydroxy, sulfhydryl, carboxy, thiocarboxy, sulfur, or an unsubstituted C₁-C₃ group or C₁-C₃ group substituted with at least one group independently selected from cyano, nitro, NH, amino, oxo, halo, hydroxy, sulfhydryl, carboxy, thiocarboxy and sulfur; or

W, V and Y form a 5 or 6 membered substituted or unsubstituted heterocyclic or carbocyclic ring fused with Z, wherein the heterocyclic ring includes from 1 to 3

~~heteroatoms selected from O, N and S, and the heterocyclic or carbocyclic ring~~, when substituted, has at least one constituent selected from cyano, nitro, NH, amino, oxo, halo, hydroxy, sulfhydryl, carboxy, thiocarboxy, sulfur, or an unsubstituted C₁-C₃ group or C₁-C₃ group substituted with at least one group independently selected from cyano, nitro, NH, amino, oxo, halo, hydroxy, sulfhydryl, carboxy, thiocarboxy and sulfur; and R is CH₂R', CXR' or CHX'R';

X is O or S;

X' is cyano, nitro, amino, halo, hydroxy, sulfhydryl, carboxy, thiocarboxy, or an unsubstituted C₁-C₃ group or C₁-C₃ group substituted with at least one group independently selected from cyano, nitro, NH, amino, oxo, halo, hydroxy, sulfhydryl, carboxy, thiocarboxy and sulfur;

R' is NH, O or S bonded to the spacer; and

Z is a phenyl ~~carbocyclic or heterocyclic~~ group, consisting of one ring independently having 5 or 6 ring members and at least two substituents independently selected from nitro, NH, amino, cyano, halo, hydroxy, carboxy, oxo, sulfur, sulfhydryl, C₁-C₂ alkoxy, C₁-C₂ acyl, or a C₁-C₂ alkyl or C₁-C₂ alkenyl group with at least one substituent selected from nitro, NH, amino, cyano, halo, hydroxy, carboxy, oxo, sulfur, sulfhydryl, C₁-C₂ alkoxy and C₁-C₂ acyl.

2-25. (Cancelled)

26. (Currently amended) The method of claim 1, ~~wherein the~~ being a method inhibits a for inhibiting dynamin-dependent endocytosis in the cell or synaptosome condition in a mammal.

27-55. (Cancelled)

56. (Currently amended) A method according to claim 26, wherein for at least one of M and M':

V is C;

W is CH; and

Y is cyano, nitro, amino, halo, hydroxy, sulfhydryl, carboxy, thiocarboxy, or an unsubstituted C₁-C₂ group or C₁-C₂ group substituted with at least one group independently selected from cyano, nitro, NH, amino, oxo, halo, hydroxy, sulfhydryl, carboxy, thiocarboxy and sulfur; or

W, V and Y form a 5 or 6 membered substituted or unsubstituted heterocyclic or carbocyclic ring fused with Z, wherein the heterocyclic ring includes from 1 to 3 heteroatoms selected from O, N and S, and the carbocyclic or heterocyclic ring, when substituted, has at least one substituent selected from cyano, nitro, NH, amino, oxo, halo, hydroxy, sulfhydryl, carboxy, thiocarboxy and sulfur, or an unsubstituted C₁-C₂ group or C₁-C₂ group substituted with at least one group independently selected from cyano, nitro, NH, amino, oxo, halo, hydroxy, sulfhydryl, carboxy, thiocarboxy and sulfur; and

R is CH₂R', CXR' or CHX'R';

X is O or S; and

X' is cyano, nitro, amino, halo, hydroxy, sulfhydryl, carboxy, thiocarboxy, or an unsubstituted C₁-C₂ group or C₁-C₂ group substituted with at least one group independently selected from cyano, nitro, NH, amino, oxo, halo, hydroxy, sulfhydryl; carboxy, thiocarboxy and sulphur.

57. (Currently amended) A method according to claim 56, wherein:

Y is cyano, nitro, amino, carboxy, hydroxy, sulfhydryl, or thiocarboxy; or

W, V and Y form a 5 or 6 membered substituted or unsubstituted heterocyclic or carbocyclic ring fused with Z, wherein the heterocyclic ring includes from 1 to 3 heteroatoms selected from O, N and S, and the carbocyclic or heterocyclic ring, when substituted, has at least one substituent selected from cyano, nitro, amino, hydroxy, sulfhydryl, carboxy and thiocarboxy, or a C₁-C₂ group substituted with a group selected from cyano, nitro, amino, hydroxy, sulfhydryl, carboxy and thiocarboxy; and

R is CXR'.

58. (Cancelled)

59. (Cancelled)

60. (Currently amended) A method according to claim ~~57~~~~[[68]]~~, wherein the Z group is an aryl group with has two of said substituents in ortho positions relative to one another.

61. (Cancelled)

62. (Currently amended) A method according to claim 60, wherein W, V and Y form ~~forms~~ a 6 membered ~~heterocyclic~~ ring fused with Z.

63. (Currently amended) A method according to claim ~~60~~~~[[64]]~~, wherein V is C and Y is cyano, nitro, amino, or ~~carboxy, hydroxy, sulphydryl or thiocarboxy.~~


64. (Cancelled)

65. (Currently amended) A method according to claim ~~60~~~~[[64]]~~, wherein the two substituents are independently selected from nitro, amino and hydroxy.

66. (Previously presented) A method according to claim 65, wherein the two substituents are hydroxy.

67-68. (Cancelled)

69. (Currently amended) A method according to claim 26, wherein M and M' are each independently a moiety as follows:



wherein X is O or S ;

Y is cyano, nitro, amino, halo, hydroxy, sulfhydryl, carboxy, or thiocarboxy; or

R₁ and Y are cyclised forming a 5 or 6 membered substituted or unsubstituted ~~heterocyclic or carbocyclic~~ ring, wherein the ~~heterocyclic ring includes 1 or 2 heteroatoms selected from O, N and S, and the carbocyclic or heterocyclic~~ ring, when substituted, has at least one substituent selected from cyano, nitro, NH, amino, oxo, halo, hydroxy, sulfhydryl, carboxy, thiocarboxy and sulfur; and


R₂ to R₅ are independently hydrogen or a substituent independently selected from nitro, amino, halo, hydroxy, carboxy, sulfhydryl, thiocarboxy, C₁-C₂ alkoxy and C₁-C₂ acyl; or

R₁ to R₅ are independently hydrogen or a substituent independently selected from nitro, amino, halo, hydroxy, carboxy, sulfhydryl, thiocarboxy, halo, C₁-C₂ alkoxy and C₁-C₂ acyl; and

R is NH, O is S bonded to the spacer Sp; and

wherein at least one of M and M' is characterised in that, at least two of R₁ to R₅ are other than hydrogen, and when R₁ to R₂ are other than hydrogen at least one of R₃ to R₅ is also other than hydrogen, or when R₁ and Y are cyclised, at least two of R₂ to R₅ are other than hydrogen.

70. (Previously presented) A method according to claim 69, wherein at least two of R_2 to R_4 are other than hydrogen.
71. (Cancelled)
72. (Previously presented) A method according to claim 70, wherein at least three of R_1 to R_4 are other than hydrogen.
73. (Currently amended) A method according to claim 72~~[[70]]~~, wherein at least two of R_2 to R_4 are hydroxy.
74. (Cancelled)
75. (Previously presented) A method according to claim 73, wherein Y is cyano, X is O and R is NH.
76. (Previously presented) A method according to claim 75, wherein M and M' are the same.
77. (Previously presented) A method according to claim 26, wherein the spacer Sp permits the compound to adopt a hairpin conformation.
78. (Previously presented) A method according to claim 26, wherein the spacer Sp comprises an unsubstituted alkane chain as follows:



wherein n is an integer of from 1 to 5.

79. (Previously presented) A method according to claim 1, wherein the compound of Formula I is a dimeric tyrphostin.

80. (Cancelled)

81. (Currently amended) A method according to claim 73, wherein X is O, R is NH and R₁ and Y are cyclised, forming a substituted ~~heterocyclic group~~ ring with 6 ring members.

82-85. (Cancelled)

86. (Currently amended) The method of claim 26, being a method for prophylaxis or treatment of ~~wherein the method prevents or treats epilepsy or inhibits a dynamin-dependent endocytosis~~ in a mammal, wherein the method ~~comprising~~ comprises administering to the mammal an effective amount of the compound of formula I, or a physiologically acceptable salt or prodrug thereof.